

Diabetes Mellitus and Blood Glucose Variability

Increase the 30-day Readmission Rate after Kidney Transplantation

Agni Orfanoudaki, PhD; Curtiss B. Cook, MD; Soroush Saghafian, PhD; Janna Castro, BS; Heidi E. Kosiosek, MS; Harini A. Chakkerla, MD, MPH

University of Oxford, Oxford, England (Dr. Orfanoudaki); Harvard Kennedy School, Harvard University, Cambridge, Massachusetts (Dr. Saghafian, Dr. Orfanoudaki); Department of Information Technology, Mayo Clinic Hospital, Phoenix, Arizona, 85054, USA (Ms Castro); Biostatistics, Mayo Clinic Hospital, Phoenix, Arizona, 85054, USA (Ms. Kosiosek); Division of Endocrinology, Mayo Clinic Arizona, Scottsdale, Arizona, 85259, USA (Dr. Cook); Division of Transplantation, Mayo Clinic Hospital, Phoenix, Arizona, 85054, USA (Dr. Chakkerla)

Corresponding Author: Harini A. Chakkerla, MD, MPH, Professor of Medicine, Division of Transplantation, Mayo Clinic Hospital, 5777 East Mayo Boulevard, Phoenix, AZ 85054.

Telephone: 480-342-0161. Fax: 480-342-2324. E-mail: chakkerla.harini@mayo.edu.

Keywords: insulin therapy, blood glucose, solid-organ transplantation, readmission risk

©2023 Mayo Foundation for Medical Education and Research

ABSTRACT

Background: Inpatient hyperglycemia, is an established independent risk factor among several patient cohorts, with hospital readmission. This has not been studied after kidney transplantation. Nearly one-third of patients who have undergone a kidney transplant reportedly experience 30-day readmission.

Methods: Data on first-time solitary kidney transplantations were retrieved between September 2015 to December 2018. Information was linked to the electronic health record to determine diagnosis of diabetes mellitus, and extract glucometric and insulin therapy data. Univariate logistic regression analysis and the XGBoost algorithm used to predict 30-day readmission. We report the average performance of the models on the testing set on five bootstrapped partitions of the data to ensure statistical significance.

Results: The cohort included 1036 patients who received kidney transplantation, 224 (22%) experienced 30-day readmission. The machine learning algorithm was able to predict 30-day readmission with an average AUC of 77.3% (95% CI 75.30-79.3%). We observed statistically significant differences in presence of pretransplant diabetes, inpatient-hyperglycemia, inpatient-hypoglycemia, minimum and maximum glucose values among those with higher 30-day readmission rates. The XGBoost model identified the index admission length of stay, presence of hyper- and hypoglycemia and recipient and donor BMI values as the most predictive risk factors of 30-day readmission. Additionally, significant variations in the therapeutic management of blood glucose by providers were observed.

Conclusions: Suboptimal glucose metrics during hospitalization after kidney transplantation is associated with an increased risk for 30-day hospital readmission. Optimizing the hospital blood glucose management, a modifiable factor, after kidney transplantation may reduce the risk of 30-day readmission.

INTRODUCTION

Early hospital readmissions are associated with increased healthcare costs and higher morbidity. Consequently, the reduction of early hospital readmissions (defined as occurring within 30 days after discharge) has become a national priority and an important quality measure.

¹ Inpatient hyperglycemia has been associated with poorer hospital outcomes, increased readmissions, increased mortality and health care costs, in a variety of populations studied, independent of a diagnosis of diabetes mellitus (DM). ²⁻⁵

In 2012, the Affordable Care Act instituted the Hospital Readmissions Reduction Program (HRRP), overseen by the Centers for Medicare and Medicaid Services (CMS). ⁶ The HRRP prioritizes the reduction of readmissions within 30 days for six conditions. ⁶ CMS is authorized to cut Medicare reimbursements to hospitals with higher than predicted risk-adjusted readmission rates for each hospital based on its performance during a rolling performance period. ⁶ Furthermore, with various efforts in measuring and publicly reporting of hospital outcomes ^{7,8} improving measures such as risk-adjusted readmission rates have become more prominent than before. Also, although not included under the HRRP umbrella, 30-day hospital readmissions are of interest for other conditions. For example, successful kidney transplantation reduces morbidity and mortality among patients with end-stage renal disease and is a cost-effective option in comparison to continued dialysis. ⁹⁻¹¹ In 2021, there were nearly 25,000 kidney transplants performed in the U.S. ¹² Studies have reported nearly one-third of patients who have undergone a kidney transplant experience readmission within 30 days. ^{13,14} Early readmissions in these patients are associated with poorer outcomes, including higher mortality, and allograft failure. ¹⁵⁻

Several studies have analyzed the variables contributing to 30-day readmission after kidney transplantation.^{16,18-20} Identifying which are modifiable could lead to changes in practice that produce better outcomes. However, although inpatient hyperglycemia has been shown to be a risk factor for hospital readmissions in various populations, the relationship between inpatient glucose control and hospital readmissions among patients receiving a kidney transplant has not been studied. We performed a retrospective analysis to examine the association between different inpatient glucometrics and risk of 30-day readmissions occurring in a cohort of patients undergoing a kidney transplant. The analysis leverages a well-established machine learning algorithm that is able to capture non-linear relationships between the risk factors and occurrence of readmissions.²¹ More broadly, our study contributes to our prior work in better understanding, measuring, screening, and managing glucometrics for patients who undergo transplantation.²²⁻²⁷

METHODS

Study Cohort

The study has been approved by Mayo Clinic Office for Human Research Protection. Patients undergoing a kidney transplant between September 25, 2015 and December 25, 2018 were retrieved from the electronic health records (EHR). Only patients undergoing first time solitary kidney transplants were selected. Individuals who required readmission within the first 30 days following the index admission were then selected. Data on age, sex, race, and length of index hospital stay were extracted. To account for differences in the complexity of care in the hospital, Medicare Severity Diagnosis Related Group (MS-DRG) values were retrieved.²⁸ International Classification of Diseases, Tenth Revision codes were utilized to determine which cases had a diagnosis of DM.

Post-transplant Immunosuppression Protocol

After a brief admission (typically hours), patients undergo kidney transplantation. As per described in our previous methodology,²² the standard immunosuppression protocol is induction therapy with either Campath (alemtuzumab), rabbit antithymocyte immunoglobulin or basiliximab. All patients receive a 5-day tapering course of glucocorticoids (methylprednisolone intravenous 500mg on day 1, 250mg on day 2, and 125mg on day 3, followed by oral prednisone 60mg on day 4 and 30mg on day 5), after which a majority of patients receive steroid-free maintenance immunosuppression using mycophenolate mofetil and tacrolimus. The few patients who receive ongoing steroid therapy (those who are immunologically higher risk or have a diagnosis of glomerulonephritis as a cause of their kidney failure) receive the same initial 5-day tapering course in an identical fashion to that given those who are on steroid avoidance and subsequently gradually lowered to a maintenance of prednisone 5mg.

Assessment of Inpatient Glycemic Control

Point-of-care blood glucose (POC-BG) measurements were used to assess glycemic control. POC-BG measurements were obtained using a NovoStat glucometer (Nova Biomedical, Waltham, MA).²⁹ POC-BG data were used to calculate the patient-stay mean glucose level ($\text{BedGluc}_{\text{avg}}$) and during the last 24-hours before discharge ($\text{L24BedGluc}_{\text{avg}}$) as has been calculated previously.^{22,23,30,31} Additional glucometrics included the average maximum ($\text{BedGlucMax}_{\text{avg}}$) and minimum ($\text{BedGlucMin}_{\text{avg}}$) POC-BG values. In accordance with current guidelines, hyperglycemia was defined as glucose >180 mg/dL.³²⁻³⁴ The percentage of patients who had a $\text{BedGluc}_{\text{avg}}$ greater than 180 mg/dL was calculated, and the proportion of patients experiencing a hypoglycemic event (POC-BG <70 mg/dL) was determined. Hemoglobin A1c (Hgb A1c) was included in the data when available.

Definitions of Inpatient Insulin Regimen

A basal-bolus regimen (e.g., long-acting insulin combined with rapid-acting insulin with meals and correction doses as needed for hyperglycemia) is the recommended method of insulin management of hyperglycemia in hospitalized patients.³²⁻³⁴ Data on administered insulin therapy was extracted from the pharmacy information system. Insulin type was classified as basal or bolus, and patterns of insulin administration were then defined as *none*, *bolus only*, or *basal-bolus* as previously described.^{30,31,35,36} Due to the small number of cases (N=7), basal-only insulin was not included. The overall regimen used during the hospital stay was determined. To examine changes in insulin regimen throughout the hospital stay, patterns of administration were determined during the first, middle, and last 24-hours (F24h, M24hr, L24h) of the hospital stay as previously described.²³ Our previous reports have shown that different insulin regimens may be applied to patients with and without DM. Thus data on insulin was stratified by DM status.

22,36

Statistical Analysis

We conducted a baseline descriptive analysis of the study cohort. Means and standard deviations (SD) were calculated for continuous variables, as well as the relative prevalence ratio in the form of percentages for categorical variables. The number of days between discharge and readmission was calculated and displayed in day increments. The distribution of insulin therapy was determined according to diabetes status. Missing data were imputed using the MedImpute algorithm, which accounts for time-dependencies in observational data.³⁷

Univariate and multivariate logistic regression models were trained to examine the association between risk conferred by various glucometrics of interest measured anytime during hospital stay as well as patient characteristics and the risk of 30-day readmission. The relative

risk ratio is reported for each independent variable as well as the corresponding 95% confidence intervals and p-values.

To enhance the power of the study we applied the XGBoost algorithm to predict the outcome of interest (30-day readmission).²¹ XGBoost is a well-established machine learning (ML) method that is commonly used in the binary classification setting for tabular data. Contrary to traditional statistical models, XGBoost can capture non-linear relationships in the data by leveraging a large number of tree-based models that are built in an iterative fashion, also known as ensemble learning. Each tree aims to improve upon the previous model. Thus, the algorithm constructs consecutive trees that correct upon the errors of earlier ones. As a result, XGBoost achieves superior performance compared to other ML algorithms. This method, also, allows us to use as input a wider set of variables that may even be highly correlated due to its ability to model non-linear interactions among the risk factors. We used as independent variables the risk factors presented in Table 1 and the metabolic treatment information for the first, middle, and last 24-hours of the admission per patient as summarized by Figures 2-3. The training of the algorithm was conducted on 75% of the total samples (in-sample partition), and 25% was used for performance evaluation (out-of-sample partition). The hyper-parameter tuning was conducted using 10-fold cross-validation. The discrimination performance of the downstream model was evaluated on 5 independent partitions of the data, to ensure statistical significance, using the Area Under the ROC Curve (AUC) metric. Each random partition maintained the same outcome prevalence ratio between the in-sample and out-of-sample datasets. To identify the risk factors that drive the algorithm's risk prediction, we computed the feature importance score, which captures the relative contribution of each variable in the derived models.³⁰ In the Results section, we report the features that ranked highest in terms of the feature importance score and directly

derive clinical insights from our machine learning method. All statistical analyses were conducted using version 3.7 of the Python programming language and the Scikit-learn library.³⁸

RESULTS

Patient Characteristics According to Readmission Status

A total of 1,036 patients underwent kidney transplant procedure. The cohort consisted of 437 patients (42%) with diabetes mellitus prior to transplant procedure. Diabetics had a longer median length of hospital stay after the procedure compared to the non-diabetics (4.0 vs 3.0, $P < 0.001$).

The highest percentage of readmissions occurred within the first 7 days following discharge for both patients with and without DM. There were 224/1,036 patients (22%) who required readmission within 30 days following discharge from their index hospital stay (Table 1). Patients requiring readmission were comparable in age, sex, and race to those not requiring readmission (Table 1). In univariate logistic regression analyses, the length of hospital stay had a small but significant association with 30-day readmission risk, rising 6% for every additional day spent in the hospital following transplant (OR 1.06; 95% CI 1.02, 1.108, $P < 0.01$). Similarly, the risk of readmission was 78% higher for every additional value of the MS-DRG index (OR 1.78; 95% CI 1.379, 2.288; $P < 0.01$) (Table 1). The risk of rehospitalization increased 97% (OR 1.97; 95% CI 1.461, 2.657; $P < 0.01$) among those with known DM.

Relationship between Inpatient Glycemic Control and Readmissions

Analysis of glycemic variables showed that patients with hyper- and hypoglycemia correlated with greater odds of readmission (Table 1). Although no significant differences were detected in $\text{BedGluc}_{\text{avg}}$ between readmission cohorts, patients who experienced at least one episode of hyperglycemia had an 86% increase in risk of 30 day readmission (OR 1.86; 95% CI

1.31,2.65; $P < 0.01$), while those who experienced hypoglycemia had a greater than 2 fold higher risk (OR 2.2; 95% CI 1.535 – 3.166; $P < 0.01$) (Table 1).

Multivariate analysis was first performed using Logistic Regression. The downstream AUC of the resulting model was 65% with (64.6%, 65.4%) 95% CI. We hypothesized that a more advanced ML algorithm, such as XGBoost, could lead to superior discrimination performance.²¹ The average AUC performance of the XGBoost models on the out of sample set for 5 independent splits of the data was 77% with (75%, 79%) 95%CI. Figure 1 illustrates the average Receiver Operator Curves for both the in-sample and out-of-sample partitions. The index admission length of stay, presence of hyper- and hypoglycemia, as well as the recipient and donor BMI values, were the 5 most predictive risk factors for the outcome of interest. Note that the XGBoost model is not a linear model, and thus, the impact of each independent variable on the outcome of interest is not fixed. Furthermore, XGBoost allows for various interactions between variables. Thus, the effect of each predictor on each individual patient changes based on the presence or absence of other risk factors. For example, the impact of hyperglycemia on a patient's risk of readmission may vary depending on the recorded BMI or the patient's age. Contrary to a logistic regression model, where the relative risk ratio does not vary for each risk factor, ML models such as XGBoost are able to capture complex relationships between the variables and generate insights more suitable for personalized medicine. For this reason, we only report the most predictive features rather than a value for each coefficient.

Patterns of Insulin Therapy

Examination of insulin regimens employed during the last 24 hours of the index hospital stay, stratified according to DM and readmission status, showed different patterns on how therapy was applied. In those DM patients not requiring readmission, overall 86% were on

insulin compared to 74% of individuals who were readmitted (Figure 2). Use of basal-bolus insulin was similar between readmission cohorts ($p=0.87$). Bolus-only insulin therapy was greater ($p<0.01$) among individuals not requiring readmission, while a greater percentage ($p<0.01$) of those readmitted were on no therapy (Figure 2). In patients without a history of DM, use of basal-bolus insulin therapy was almost never ($N=3$) applied, and the majority were on bolus-only insulin therapy during the last 24 hours of the hospital stay. The proportion of patients on bolus-only or no insulin therapy was similar between readmission categories (Figure 2).

There was a lack of consistent pattern of how insulin therapy was utilized throughout the course of the hospital stay. It appeared very varied and based on the discretion of the provider. For instance, among patients with DM (Figure 3, top panel), some patients would remain on basal-bolus or bolus only therapy throughout the hospital, while in other instances, the type of insulin therapy would vary from time segment to time segment. Similar observations were noted for those without DM (Figure 3, bottom panel).

DISCUSSION

It is well established that hospital readmission is associated with increased morbidity, mortality, and cost of care. It is also a key metric employed by CMS and other organizations to rate the performance of hospitals and healthcare systems. With the increasing efforts in improving the transparency of the outcomes in the healthcare sector through public reporting and other activities,^{7,8} reducing readmissions has become more important than ever. Suboptimal glucose metrics during hospitalization are among the factors known to be associated with an increase in readmissions in various disease states.²⁻⁵ This analysis is among the few studies to evaluate the relationship between such suboptimal glucose metrics and hospital readmissions among patients receiving a kidney transplant.³⁹⁻⁴¹ We report that the presence of hypoglycemia

and hyperglycemia during the hospitalization after kidney transplantation as well as the recipient and donor BMI values are important risk factors for 30-day hospital readmission. Our results are derived via a machine learning model that achieves good discrimination performance (AUC) on five bootstrapped samples of unseen data. Consistent with prior reports from the authors' institution, most patients with DM were on basal-bolus insulin the 24-hours prior to discharge. Insulin therapy was also deemed necessary among a large percentage of patients without DM, although a bolus-only regimen was felt to be adequate to achieve glucose control. A more detailed assessment of how insulin regimens were used showed a lack of consistency in how therapies were applied throughout the hospital stay. This "therapeutic chaos" with regards to the use of inpatient insulin may be due to the changing clinical circumstances of the patient (e.g., tapering of steroids or change in nutritional intake), or from a lack of understanding on when to use specific regimens. Most likely it is a combination of factors. Practitioners tend to make decisions about insulin in a reactive rather than in a predictive manner, i.e., they make changes based on the previous day's glucose patterns. In the absence of systematic clinical guidelines and due to the dynamic and complex trajectory of glucose metrics, there is an increased risk of uncontrolled hyper- and hypoglycemia. There are a few reported studies describing a personalized approach in diabetes management after kidney transplant. Our study highlights the need for robust studies to be done to study and prescribe the approach in diabetes management after a kidney transplant. Specifically, future studies could focus on the personalization of the optimal in-hospital insulin therapy after kidney transplantation based on a broader set of patient characteristics that dynamically change during the hospital stay. Future research can also extend our analysis to other major organ transplantations and contribute to the stream of the literature

that aims at findings similarities and differences between kidney and other solid organ transplant patients.^{24,25}

Limitations

Our study has a few limitations including this was a single academic medical center and analysis used retrospective data from EHRs. As a result, other confounding factors may exist that could have affected the patients' trajectory with glucose control which is not included in the data including setting of infection, presence of delayed graft function and overall a sicker cohort of patients. The difference in the performance of the ML model between the in-sample and out-of-sample data partitions is attributed to overfitting and could be remedied in future studies by expanding the sample size of the population. This retrospective analysis does not provide insight into the reasons underlying the decision-making behavior of clinicians regarding clinical assessment and prescriptions. This pertains to the inability to discover causal relations between the variables and the outcome of interest, which is not the output of our findings even though there is high degree of association connectivity between the two. In addition, blood glucose values were not captured in the form of continuous variables; instead, we defined three periods during which the blood glucose and insulin usage were recorded (first 24 h, middle 24 h, and the last 24 h). However, since the median length of hospital stay was three days, we expect that the proposed time periods capture all necessary information regarding metabolic factors.

CONCLUSIONS

Our results are among the first few reported studies studying the consequences of uncontrolled in-hospital blood glucose measurements after a kidney transplant, we demonstrated a higher risk of 30-day readmission. Our findings highlight the need for further investigation of

metabolic factors and their association with the risk of hospital readmission across other transplanted organs, as well as novel strategies to manage blood glucose measurements. We believe that medical care institutions would benefit from standardization of protocols that optimize the in-hospital blood glucose management in this patient population which may minimize the likelihood of 30-day readmission and decreasing healthcare costs and improve allograft and patient survival.

Disclosures: The authors declare no conflicts of interest.

Funding: None

Personal Thanks: Barbara Weisser from the Academic Support Office, Mayo Clinic, Scottsdale, Arizona for her assistance in the preparation of the manuscript.

Author Contributions: A.O. conducted the analysis, contributed to the discussion, wrote, reviewed, and edited the manuscript. J.C. and H.K. collected the data. H.C. and C.C. led the data collection, contributed to the discussion and results interpretation, wrote, reviewed, and edited the manuscript. S.S. reviewed and edited the manuscript. All authors approved the final version of the manuscript. A.O. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Preliminary data was submitted and accepted as a poster for the 2022 American Transplant Congress meeting.

Data Sharing Statement: The data underlying this article will be shared on reasonable request to the corresponding author.

References

1. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med*. Apr 2 2009;360(14):1418-28.
doi:10.1056/NEJMsa0803563
2. Umpierrez GE, Isaacs SD, Bazargan N, You X, Thaler LM, Kitabchi AE. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab*. Mar 2002;87(3):978-82. doi:10.1210/jcem.87.3.8341
3. Williams LS, Rotich J, Qi R, et al. Effects of admission hyperglycemia on mortality and costs in acute ischemic stroke. *Neurology*. Jul 9 2002;59(1):67-71. doi:10.1212/wnl.59.1.67
4. Foo K, Cooper J, Deaner A, et al. A single serum glucose measurement predicts adverse outcomes across the whole range of acute coronary syndromes. *Heart*. May 2003;89(5):512-6.
doi:10.1136/heart.89.5.512
5. Whitcomb BW, Pradhan EK, Pittas AG, Roghmann MC, Perencevich EN. Impact of admission hyperglycemia on hospital mortality in various intensive care unit populations. *Crit Care Med*. Dec 2005;33(12):2772-7. doi:10.1097/01.ccm.0000189741.44071.25
6. Centers for Medicare and Medicaid Services. Readmissions Reduction Program. Available at: <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/Readmissions-Reduction-Program> Accessed January 7, 2022.
2012;
7. Saghafian S, Hopp WJ. Can public reporting cure healthcare? The role of quality transparency in improving patient-provider alignment. *Operations Research*. 2020;68(1):71-92.

8. Saghafian S, Hopp WJ. The Role of Quality Transparency in Health Care: Challenges and Potential Solutions. *NAM Perspect.* 2019;2019doi:10.31478/201911a
9. Axelrod DA, Schnitzler MA, Xiao H, et al. An economic assessment of contemporary kidney transplant practice. *Am J Transplant.* May 2018;18(5):1168-1176. doi:10.1111/ajt.14702
10. Wolfe RA, McCullough KP, Leichtman AB. Predictability of survival models for waiting list and transplant patients: calculating LYFT. *Am J Transplant.* Jul 2009;9(7):1523-7. doi:10.1111/j.1600-6143.2009.02708.x
11. Wolfe RA, Ashby VB, Milford EL, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med.* Dec 2 1999;341(23):1725-30. doi:10.1056/NEJM199912023412303
12. U.S. Department of Health & Human Services. Organ Procurement and Transplantation Network. Available at <https://optn.transplants.hrsa.gov/data/view-data-reports/build-advanced/> Accessed December 7, 2021.
13. Lubetzky M, Yaffe H, Chen C, Ali H, Kayler LK. Early Readmission After Kidney Transplantation: Examination of Discharge-Level Factors. *Transplantation.* May 2016;100(5):1079-85. doi:10.1097/TP.0000000000001089
14. McAdams-DeMarco MA, Law A, Salter ML, et al. Frailty and early hospital readmission after kidney transplantation. *Am J Transplant.* Aug 2013;13(8):2091-5. doi:10.1111/ajt.12300
15. Li AH, Lam NN, Naylor KL, Garg AX, Knoll GA, Kim SJ. Early Hospital Readmissions After Transplantation: Burden, Causes, and Consequences. *Transplantation.* Apr 2016;100(4):713-8. doi:10.1097/TP.0000000000000917

16. Harhay M, Lin E, Pai A, et al. Early rehospitalization after kidney transplantation: assessing preventability and prognosis. *Am J Transplant*. Dec 2013;13(12):3164-72.
doi:10.1111/ajt.12513
17. McAdams-Demarco MA, Grams ME, King E, Desai NM, Segev DL. Sequelae of early hospital readmission after kidney transplantation. *Am J Transplant*. Feb 2014;14(2):397-403.
doi:10.1111/ajt.12563
18. Famure O, Kim ED, Au M, et al. What Are the Burden, Causes, and Costs of Early Hospital Readmissions After Kidney Transplantation? *Prog Transplant*. Jun 2021;31(2):160-167. doi:10.1177/15269248211003563
19. McAdams-Demarco MA, Grams ME, Hall EC, Coresh J, Segev DL. Early hospital readmission after kidney transplantation: patient and center-level associations. *Am J Transplant*. Dec 2012;12(12):3283-8. doi:10.1111/j.1600-6143.2012.04285.x
20. Hogan J, Arenson MD, Adhikary SM, et al. Assessing Predictors of Early and Late Hospital Readmission After Kidney Transplantation. *Transplant Direct*. Aug 2019;5(8):e479.
doi:10.1097/TXD.0000000000000918
21. Chen T, Guestrin C. XGBoost: A scalable tree boosting system. presented at: KDD '16: Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining; 2016;
22. Chakkerla HA, Weil EJ, Castro J, et al. Hyperglycemia during the immediate period after kidney transplantation. *Clin J Am Soc Nephrol*. Apr 2009;4(4):853-9.
doi:10.2215/CJN.05471008

23. Munshi VN, Saghafian S, Cook CB, Steidley DE, Hardaway B, Chakkera HA. Incidence, Risk Factors, and Trends for Postheart Transplantation Diabetes Mellitus. *Am J Cardiol*. Feb 1 2020;125(3):436-440. doi:10.1016/j.amjcard.2019.10.054
24. Munshi VN, Saghafian S, Cook CB, Aradhyula SV, Chakkera HA. Use of Imputation and Decision Modeling to Improve Diagnosis and Management of Patients at Risk for New-Onset Diabetes After Transplantation. *Ann Transplant*. Mar 16 2021;26:e928624. doi:10.12659/AOT.928624
25. Munshi VN, Saghafian S, Cook CB, Werner KT, Chakkera HA. Comparison of post-transplantation diabetes mellitus incidence and risk factors between kidney and liver transplantation patients. *PLoS One*. 2020;15(1):e0226873. doi:10.1371/journal.pone.0226873
26. Bolori A, Saghafian S, Chakkera HA, Cook CB. Characterization of Remitting and Relapsing Hyperglycemia in Post-Renal-Transplant Recipients. *PLoS One*. 2015;10(11):e0142363. doi:10.1371/journal.pone.0142363
27. Bolori A, Saghafian S, Chakkera HA, Cook CB. Data-driven management of post-transplant medications: an ambiguous partially observable Markov decision process approach. *Manufacturing and Service Operations Management*. 2020;22(5):1066-1087.
28. Centers for Medicare and Medicaid Services. MS-DRG Classifications and Software. Available at <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/MS-DRG-Classifications-and-Software> Accessed January 7, 2022.
29. Leighton ME, Thompson BM, Castro JC, Cook CB. Nurse adherence to post-hypoglycemic event monitoring for hospitalized patients with diabetes mellitus. *Appl Nurs Res*. Dec 2020;56:151338. doi:10.1016/j.apnr.2020.151338

30. Cook CB, Castro JC, Schmidt RE, et al. Diabetes care in hospitalized noncritically ill patients: More evidence for clinical inertia and negative therapeutic momentum. *J Hosp Med.* Jul 2007;2(4):203-11. doi:10.1002/jhm.188
31. Mackey PA, Boyle ME, Walo PM, Castro JC, Cheng MR, Cook CB. Care directed by a specialty-trained nurse practitioner or physician assistant can overcome clinical inertia in management of inpatient diabetes. *Endocr Pract.* Feb 2014;20(2):112-9. doi:10.4158/EP13201.OR
32. Umpierrez GE, Hellman R, Korytkowski MT, et al. Management of hyperglycemia in hospitalized patients in non-critical care setting: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab.* Jan 2012;97(1):16-38. doi:10.1210/jc.2011-2098
33. Society of Hospital Medicine. *The Glycemic Control Implementation Guide: Improving Glycemic Control, Preventing Hypoglycemia and Optimizing Care of the Inpatient with Hyperglycemia and Diabetes.* Available at <https://www.hospitalmedicine.org/globalassets/clinical-topics/clinical-pdf/gcmi-guide-m4.pdf>. 2015.
34. American Diabetes Association Professional Practice Committee, Draznin B, Aroda VR, et al. Diabetes Care in the Hospital: Standards of Medical Care in Diabetes-2022. *Diabetes Care.* Jan 1 2022;45(Suppl 1):S244-S253. doi:10.2337/dc22-S016
35. Coan KE, Schlinkert AB, Beck BR, et al. Clinical inertia during postoperative management of diabetes mellitus: relationship between hyperglycemia and insulin therapy intensification. *J Diabetes Sci Technol.* Jul 1 2013;7(4):880-7. doi:10.1177/193229681300700410

36. Werner KT, Mackey PA, Castro JC, Carey EJ, Chakkerla HA, Cook CB. Hyperglycemia during the immediate period following liver transplantation. *Future Sci OA*. Mar 2016;2(1):FSO97. doi:10.4155/fsoa-2015-0010
37. Bertsimas D, Orfanoudaki A, Pawlowski C. Imputation of clinical covariates in time series. *Mach Learn*. 2021;110:185-248. doi:https://doi.org/10.1007/s10994-020-05923-2
38. Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine learning in python. *J Mach Learn Research*. 2011;12(85):2825-2830.
39. Ramirez SC, Maaske J, Kim Y, et al. The association between glycemic control and clinical outcomes after kidney transplantation. *Endocr Pract*. Sep 2014;20(9):894-900. doi:10.4158/EP13463.OR
40. Hosseini MS, Nemati E, Pourfarziani V, Taheri S, Nourbala MH, Einollahi B. Early hyperglycemia after allogenic kidney transplantation: does it induce infections. *Ann Transplant*. 2007;12(4):23-6.
41. Orfanoudaki A, Saghafian S, Song K, Chakkerla H, Cook C. Algorithm, Human, or the Centaur: How to Enhance Clinical Care? Available at: SSRN 4302002. 2022;

Table

Table 1: Comparison of patients undergoing first time kidney transplant (September 2015 to December 2018), according to 30-day readmission status.*

30-day readmission status	Yes N=224	No N=812	OR (CI)	P-values
Age, y	53(14)	53(14)	1.0 (0.99,1.01)	0.933
Male sex (%)	60	59	1.04 (0.77,1.40)	0.808
White race (%)	66	67	0.93 (0.68,1.28)	0.666
Length of index hospital stay, d	5.1(3.8)	4.1(3.6)	1.06 (1.02,1.11)	<0.01
MS-DRG	3.52(0.71)	3.35(0.41)	1.78 (1.38,2.29)	<0.001
Donor Deceased (%)	87	78	1.58 (1.01,2.46)	0.045
Diabetes, (%)	55	39	1.97 (1.46,2.66)	<0.001
Glucometrics				
BedGluc _{avg}	152(32)	140(30)	1.0 (0.99,1.01)	0.430
Patients with hyperglycemia (At least one value of BedGluc >180 mg/dL, (%))	79	66	1.86 (1.31,2.65)	<0.001
Patients with hyperglycemia (BedGluc _{avg} >180 mg/dL, (%))	17	15	1.19 (0.80,1.78)	0.390
Patients with hypoglycemia (%)	25	13	2.2 (1.54,3.17)	<0.001
BedGlucMax _{avg}	245(79)	224(71)	1.0 (1.00,1.01)	<0.001
BedGlucMin _{avg}	87(25)	94(24)	0.99 (0.98,1.00)	<0.001

*Data are medians (± SD) for continuous variables

LOS (length of stay); MSDRG (Medicare Severity-Diagnosis Related Group); KDPI (Kidney Donor Profile Index) BedGluc_{avg} (patient stay mean glucose); L24BedGluc_{avg} (average glucose 24 hours prior to discharge); BedGlucMax_{avg} (average maximum glucose); BedGlucMin_{avg} (average minimum glucose)

Figure Legends

Figure 1: Area under the Receiver Operator Curves for the in sample (training) and out of sample (testing) data.

Figure 2: Insulin Regimen 24 Hours Prior to Discharge.

Figure 3. Distribution of insulin treatment regimens during the hospital admission. The percentages indicate the proportion of the population that followed each trajectory.

Figure 1




Figure 1: Area under the Receiver Operator Curves for the XGBoost algorithm for the in-sample (training) and out-of-sample (testing) data.

Figure 2




Figure 2: Insulin Regimen 24 Hours Prior to Discharge.

Figure 3




Figure 3. Distribution of insulin treatment regimens during the hospital admission. The percentages indicate the proportion of the population that followed each trajectory